


For Reference

NOT TO BE TAKEN FROM THIS ROOM

Ex libris
UNIVERSITATIS
ALBERTAE NSIS





Digitized by the Internet Archive
in 2022 with funding from
University of Alberta Library

<https://archive.org/details/Bates1984>

T H E U N I V E R S I T Y O F A L B E R T A

RELEASE FORM

NAME OF AUTHOR Larry Bates

TITLE OF THESIS HIDDEN SYMMETRY IN MONOPOLES

DEGREE FOR WHICH WAS PRESENTED M.Sc.

YEAR THIS DEGREE GRANTED 1984

Permission is hereby granted to THE UNIVERSITY OF ALBERTA LIBRARY to reproduce single copies of this thesis and to lend or sell such copies for private, scholarly or scientific research purposes only.

The author reserves other publication rights, and neither the thesis nor extensive extracts from it may be printed or otherwise reproduced without the author's written permission.

THE UNIVERSITY OF ALBERTA

HIDDEN SYMMETRY IN MONOPOLES

by



Larry Bates

A THESIS

SUBMITTED TO THE FACULTY OF GRADUATE STUDIES AND RESEARCH
IN PARTIAL FULFILLMENT OF THE REQUIREMENTS FOR THE DEGREE OF
MASTER OF SCIENCE

DEPARTMENT OF MATHEMATICS

EDMONTON, ALBERTA

FALL 1984

THE UNIVERSITY OF ALBERTA

FACULTY OF GRADUATE STUDIES AND RESEARCH

The undersigned certify that they have read, and recommend to the
Faculty of Graduate Studies and Research, for acceptance, a thesis
entitled

HIDDEN SYMMETRY IN MONOPOLES

submitted by Larry Bates

in partial fulfillment of the requirements for the degree of
Master of Science

ABSTRACT

The motion of a classical charged particle in the field of a magnetic monopole with a specially chosen potential is studied from a symplectic-geometric point of view. Throughout we emphasize the global structure of the space of solutions and its symmetries. In particular, a systematic geometric algorithm is presented that constructs the first integrals of the motion.

ACKNOWLEDGEMENTS

To Hans Künzle, who opened the door to that most beautiful and charming of mathematical rooms, the symplectic universe. Your genial manner and felicitous remarks during many long discussions on monopoles, geometry or whatever helped me no end. The fact that this thesis is finished is due in large part to your never-failing interest and patience.

To Jim Nester, for your many hours of listening and good advice.

To the other graduate students in mathematics, especially Brian, Joseph and Ron, for your help and encouragement.

Not all debts incurred are mathematical, and to Mary, Brett, David and Gordon, my deepest thanks for your support.

To Christine and Laura for their first-rate typing and coping with two frustrating word processors, one of them the author of this thesis.

INTRODUCTION

A magnetic monopole is a magnetic charge. That is to say, a magnetic field that is spherically symmetric and falls off as the inverse of the square of the distance from the charge. Despite have never been found in nature, magnetic monopoles form a central part of the current understanding of theoretical physics, running the gamut of gauge theories to elementary particles to cosmology. This popularity is due in large measure to the insight gained into current theories by sophisticated but solvable examples. For some flavour of current work the reader is referred to Balachandran et.al.(1980), Barut et.al.(1971), Carrigan (1965), D'Hoker and Vinet (1983-1984), Hitchin (1982), Jaffe and Taubes (1980), Miller (1976), Moriyasu (1981), Sanders (1966), T'Hooft (1974), and Wu and Yang (1975-1976).

In this essay we add a specially chosen potential to this magnetic field and study the motion of a classical charged particle in this field (see Boulware et.al.(1976), Schonfeld (1980), and Zwanziger (1968)). This provides a one-parameter family of inequivalent Hamiltonian systems that all "look somewhat like" the Kepler problem (for brevity we now refer to this system as the monopole problem).

In the prolegomenon we give some introductory remarks on the equations of motion and then, following Abraham and Marsden, introduce the notion of a Hamiltonian system to give a global construction of the monopole problem. The reader having little sympathy with bundle

constructions and momentum mappings, that is to say the current vogue in mechanics, can proceed directly to chapter two. In defense of this construction, which will seem incredibly abstract and obfuscating, it seems only right to point out that this is the only approach I know of that not only does the construction globally, but preserves the physical interpretation of variables. We provide enough detail to correct a statement of Marsden (1981) on the introduction of charge in this construction.

In chapter one we develop a geometric algorithm based on complete lifts that finds conserved quantities of the motion. In particular, we find the (corrected cf. Schonfeld) generators of a symplectic $SO(4)$ action on negative energy orbits. A by-product of this symplectic approach is the explanation of why the monopole problem is Kepler-like. In this chapter, sections two and three are essentially original material.

Chapter two exploits techniques due to Souriau (1974) to integrate the equations of motion and determine the topology of energy surfaces. The idea that these techniques could be extended to the monopole problem is due to Hans Künzle.

Finally, in the eschata we collect some useful definitions, fix notation, and explain some notions from mechanics and geometry.

PROLEGOMENON

1. THE MODEL

A magnetic monopole has a magnetic field \vec{B} , with

$$\vec{B} = \frac{g}{r^3} \vec{r} \quad \vec{r} \in \mathbb{R}^3, \quad r = \|\vec{r}\|.$$

Now consider the equation of motion of a classical charged particle of charge e in the field of a magnetic monopole (with specially chosen potential and setting $\lambda = eg$)

$$m\ddot{\vec{r}} = \frac{\lambda \vec{r} \times \dot{\vec{r}}}{r^3} - \nabla \left(-\frac{k}{r} + \frac{\lambda^2}{2mr^2} \right). \quad (1)$$

Note that when $\lambda = 0$ we have the Kepler problem. Now in (1) we define

$$\vec{J} = m\vec{r} \times \dot{\vec{r}} + \lambda \hat{r}, \quad \hat{r} = \frac{1}{r} \vec{r}$$

and calculate that $\dot{\vec{J}} = 0$.

Now

$$\hat{r} \cdot \vec{J} = \lambda$$

so the motion of the particle lies on a cone of pitch α with

$$\cos \alpha = \frac{\lambda}{J} \quad , \quad J = ||\vec{J}||$$

Now define another vector \vec{R} by

$$\vec{R} = \frac{1}{\sin \alpha} [\vec{r} - \hat{J}(\vec{r} \cdot \hat{J})] \quad .$$

Then

$$R = ||\vec{R}|| = r$$

$$\vec{R} \cdot \vec{J} = 0$$

$$m\vec{R} \times \dot{\vec{R}} = \vec{J} \quad (1a)$$

$$\ddot{\vec{R}} = -\nabla \left(\frac{-k}{R} \right)$$

so we have transformed (1) into the Kepler problem!

At this stage one might be tempted to conjecture that the reason this works is that if we rewrote (1) in Hamiltonian form, then there is a canonical transformation between (1) and the Kepler problem.

A local construction of a Lagrangian can be done if we find a potential

$$A = A_k dx^k \quad \text{for the electromagnetic field tensor} \quad F = \frac{1}{2} F_{ij} dx^i \wedge dx^j ,$$

where $F = dA$.

Now $F_{ij} = \epsilon_{ijk} B^k$ and so

$$F = \frac{1}{2} \frac{g}{r} \epsilon_{ijk} x^i dx^j \wedge dx^k \quad \text{in cartesian coordinates}$$

$$= g \sin \phi \, d\theta \wedge d\phi \quad \text{in polar coordinates (see appendix).}$$

So, by inspection we may choose

$$A = g \cos \phi \, d\theta ,$$

and the Lagrangian has the form

$$L = \frac{1}{2} m \, v^2 + eA \cdot v + V(x)$$

and so the Hamiltonian is

$$\begin{aligned} H &= A \cdot v - L \\ &= \frac{1}{2m} (\hat{p} - eA)^2 + V(x) \end{aligned} \tag{2}$$

where \hat{p} is the canonical momentum and the equations of motion are just Hamilton's equations. (see Crampin 1981, Havas 1957).

There is a serious objection to this procedure, however. The construction of the Lagrangian and Hamiltonian are only local, while the second order equation is globally well defined. In fact, one can prove that there is no global Lagrangian for this second-order equation. To remedy this we introduce the notion of a Hamiltonian system: (the basic geometrical ideas are well-covered in Abraham-Marsden, (1978) Souriau, (1970), Arnold (1978), Maclane, (1970)).

A Hamiltonian system is the triple (P, ω, h) where P is a manifold, ω is a symplectic form on P , and h is a function called the Hamiltonian. The equation of motion is given by $i_{X_h} \omega = dh$, where X_h is the symplectic gradient of h . From this geometric viewpoint we can formulate the monopole problem globally as

$$P = (\mathbb{R}^3 \setminus \{0\}) \times \mathbb{R}^3$$

$$\omega = \frac{1}{2} \frac{\lambda}{r^3} \varepsilon_{ijk} x^i dx^j \wedge dx^k + dx^k \wedge dp_k. \quad (3)$$

$$h = \frac{p^2}{2m} + \frac{\lambda^2}{2mr^2} - \frac{k}{r}.$$

where $p_k = m\dot{x}^k$.

On the other hand, the Kepler problem is

$$\bar{P} = P$$

$$\Omega = dx^k \wedge dp_k \quad (4)$$

$$H = \frac{p^2}{2m} - \frac{k}{r}.$$

The first result that may seem initially surprising in view of (1a) is that there is no canonical transformation between (3) and (4). In fact, a careful examination of the proof of this shows that the systems (3) are not equivalent by canonical transformation for any two distinct values of λ .

We thus have an interesting example of a one-parameter family of inequivalent Hamiltonian systems that all "look somewhat like" the Kepler problem. This is explained in I.2 using the geometrical reduction scheme of Marsden and Weinstein (This is the precise way of saying the classical notion of "freezing out the angular variables"). For more background and technical details one should consult Abraham-Marsden (1978), Marsden-Weinstein (1974), Giachetti (1981), Iwai (1980); and Marmo et.al. (1979).

It is legitimate to ask why the term $\frac{\lambda^2}{2mr^2}$ occurs in the potential in (1) . The reason is that this is precisely the condition we need to give us a conserved Lenz-like vector, just as in the Kepler problem. Furthermore, it again generates an action of $SO(4)$ on our phase space by canonical transformations.

The rest of this section is devoted to explaining why the magnetic monopole problem with potential (hereafter the monopole) can be written globally as the Hamiltonian system (3) .

This is probably the most difficult section of the thesis, and, unless the reader wishes to see this, should probably proceed to the notes, keeping in mind that this construction is to show why the addition of a magnetic fields modifies the geometry of phase space (that is, the symplectic structure) but does not change the energy or our concept of momentum. The geometry is altered because now the velocities do not Poisson commute.

This construction is based on Sternberg (1977), Weinstein (1978) and Guillemin and Sternberg (1978). See also Marsden (1981). The relevant background on principal bundles and connections can be found in Choquet-Bruhat (1977), Kobayashi-Nomizu (1963), Poor (1981), Sternberg (1963), or Spivak (1974). For connections with monopoles see Cant (1981). Somewhat different approaches are in Duval and Horvathy (1982) and Montgomery (1983).

The current model for electromagnetism is a principal bundle M over space or space-time Q with structure group $G = U(1) = SO(2)$, with the electromagnetic field tensor F occurring as the curvature form of a connection on M . In this thesis we are interested in non-relativistic mechanics so we treat Q as space. That is to say, $Q = \mathbb{R}^3 \setminus \{0\}$. We also treat the connection on M as representing a vector potential for the magnetic field. We now want to build an associated symplectic manifold P , with a Hamiltonian, that is the Hamiltonian system for the monopole.

We choose local coordinates on the cotangent bundle of M , T^*M , as (q, α, z, ζ) (a fibre chart) where

$$q \in Q$$

$$\alpha \in G$$

$$z \in \mathbb{R}^{3*}$$

$$\zeta \in \mathfrak{g}^*, \text{ where } \mathfrak{g} \text{ is the Lie algebra of } G (\mathfrak{g} \approx \mathbb{R})$$

The canonical forms on T^*M have the local expression

$$\theta_M = z_k dq^k + \zeta d\alpha$$

$$\omega_M = dq^k \wedge dz_k + d\alpha \wedge d\zeta.$$

The right action of G on M lifts to a symplectic right action on T^*M with the momentum map (see appendix)

$$J_M: T^*M \rightarrow \mathfrak{g}^*: (q, \alpha, z, \zeta) \mapsto \zeta$$

To this right action we may associate the left action whose momentum map is $-J_M$. Now consider the Hamiltonian G space (see Abraham-Marsden (1978) p. 276) $S = T^*G$ with the left G action and canonical symplectic structure. The momentum map on S is

$J_G: S \rightarrow \mathfrak{g}^*: (\delta, e) \mapsto (e)$ in a local fibre chart. Now the momentum map J on $T^*M \times S$ is just $J = -J_M + J_G$. We construct the reduced manifold $P = (T^*M \times S)_0$ as $P = J^{-1}(0)/G_0$ where G_0 is the isotropy group of $0: G_0 = \{\alpha \in G \mid \text{Ad}_{\alpha}^{-1}(0) = 0\} = G$.

Thus we have an inclusion map i :

$$i: P \rightarrow T^*M \times S$$

which has the local form

$$(q, p) \mapsto (q, 0, p, -e, 0, e)$$

Note that now $\{e\}$ is a co-adjoint orbit of G in \mathfrak{g}^* , which we will identify as the charge of the particle. The symplectic form $\omega_M + \omega_G$ on $T^*M \times S$ pulls back via i to give the induced form $i^*(\omega_M + \omega_G) = \omega_P$ on P which has the local form $dq^k \wedge dp_k$.

To get a map from P to T^*Q we pick a connection on M . In a local chart for TM we have $(q^k, \alpha, \partial_k, \partial_\alpha)$ while a local chart for TQ is (q^k, ∂_k) . A connection is a way of specifying a horizontal subspace of $T_{(q, \alpha)}M$ which we define by the map $\gamma: T_q Q \rightarrow T_{(q, \alpha)}M$

$\partial_k \rightarrow \partial_k + A_k \partial_\alpha$
 whose dual $\gamma_q^*: T_{(q,\alpha)}^* M \rightarrow T_q^* Q$ takes

$$dq^k \rightarrow dq^k$$

$$d\alpha \rightarrow A_k dq^k.$$

Putting all the cotangent spaces together we get a map

$$\gamma^*: T^* M \rightarrow T^* Q: (q, \alpha, z, \zeta) \rightarrow (q, y)$$

where $y_k = z_k + A_k \zeta$.

Given a Hamiltonian H on $T^* Q$ we now pull it back to P to get a Hamiltonian $H_\gamma = h = \gamma^*(H)$. On $T^* Q$ we set $H = \frac{1}{2m} y^2 + V(q)$ and so h has the local form

$$h = \frac{1}{2m} (p - eA)^2 + V(q).$$

Making the transformation $p \rightarrow p + eA$ we find that we can write the system globally as

$$h = \frac{1}{2m} p^2 + V(x)$$

$$\omega = \frac{1}{2} \frac{\lambda}{r^3} \varepsilon_{ijk} q^i dq^j \wedge dq^k + dq^k \wedge dp_k$$

where (q, p) are cartesian coordinates on $P \approx T^* Q$.

2. NOTES:

(A) It is also interesting to see why the potential V was chosen the way it was. The reason is primarily aesthetic. It turns out that this is the condition we need so that we have an $SO(4)$ symmetry of the Hamiltonian. This potential does not come from any field theory that I am aware of, although a potential differing by a factor of two in the $\frac{1}{r^2}$ term does come from the Newtonian theory of Kunzle (1972) as a non-relativistic limit of the Newton-Cartan equations in Taub-Nut space. Further discussion is in Schonfeld (1980).

(B) We now want to look at the cohomology class of ω and note some interesting consequences and implications. We set $\hat{\omega} = \omega + d\theta_0$, so we clearly have that $\hat{\omega}$ and ω are cohomologous. Furthermore, $\hat{\omega}$ can be thought of as a form on Q instead of $P = T^*Q$. Now S^2 , the unit sphere in three-space, is a deformation retract of Q so we may evaluate the cohomology class of $\hat{\omega}$ on S^2 . This induced form on S^2 is just the restriction of $\hat{\omega}$ since there are no terms involving dr in $\hat{\omega}$. Thus the cohomology class of ω , call it $[\omega]$, is

$$[\omega] = \lambda \int_{S^2} \sin \phi \, d\phi \, d\theta$$

$$= 4\pi\lambda.$$

This further implies that there is no global formulation of the monopole problem as a Lagrangian system. This is in some sense the ultimate justification for the seemingly abstract approach to formulating the equations of motion. It is well-known in geometric quantization that for a classical problem to be quantizable the cohomology class of the symplectic structure must be an integral multiple of Planck's constant. In other words

$$[\frac{1}{h} \omega] \in H^2(P, \mathbb{Z}).$$

This only happens if the charge λ satisfies

$$\lambda = \frac{n\hbar}{2},$$

which is exactly the quantization condition of Dirac (1931).

CHAPTER ONE

We now proceed to systematically exploit the symmetry of the monopole with a symplectic methodology. In this way we avoid the rather ad hoc and 'coincidental' approaches of Boulware, Schonfeld, and Zwanziger, because the symmetry is then seen as naturally reflecting the intrinsic geometry of the system.

1. THE OBVIOUS SYMMETRY GROUP.

An obvious symmetry is an infinitesimal symplectomorphism generated by a vector field on P that

(OS1) is the complete lift of a vector field on Q ,

(OS2) is tangent to an energy surface.

In the tangent formulation this condition reads as

$$L_{\hat{\chi}} \omega = 0, \tag{1}$$

where $\hat{\chi}$ is the complete lift of χ . In other words, if

$$\chi = f^k \partial_k,$$

then $\hat{\chi} = f^k \partial_k + v^\ell \partial_\ell f^k \partial_k$. Now $L_{\hat{\chi}} \omega = \iota_{\hat{\chi}} d\omega + d\iota_{\hat{\chi}} \omega = d\iota_{\hat{\chi}} \omega$ since ω is closed.

Substituting, we get

$$L_{\tilde{\chi}} \omega = \partial_{(j} f_{k)} dq^j \wedge dv^k + \partial_a (f^k_{\omega_{kb}} - v^\ell \partial_\ell f_b) dq^a \wedge dq^b.$$

To satisfy the condition $L_{\tilde{\chi}} \omega = 0$ we must have that the individual coefficients vanish. We now need the following:

Lemma: The condition $\partial_{(j} f_{k)} = 0$ implies f is an isometry.

Proof: Suppose we have $Df + Df^T = 0$ for all u and v , that is

$$\langle Df(u), v \rangle + \langle u, Df(v) \rangle = 0.$$

By differentiating both sides, we get

$$\langle D^2 f(u, w), v \rangle + \langle u, D^2 f(v, w) \rangle = 0 \quad \text{for all } u, v, w.$$

By playing with these conditions, we derive that

$$\begin{aligned} \langle D^2 f(u, w), v \rangle &= -\langle u, D^2 f(v, w) \rangle \\ &= -\langle D^2 f(w, v), u \rangle \end{aligned}$$

$$= \langle D^2 f(v, u), w \rangle$$

$$= -\langle D^2 f(u, w), v \rangle.$$

Thus $D^2 f(u, v) = 0$ for all u and v . Therefore, we conclude that

$$f = A \cdot q + B,$$

where $A + A^T = 0$.

The other terms give no further information. Inspection of the condition $L_{\hat{X}} E = 0$ shows that B is zero. This means that f is an element of $\mathfrak{so}(3)$, the Lie algebra of $SO(3)$.

We conclude that the obvious symmetry group of the monopole is $SO(3)$, with its standard linear action.

2. THE MOMENTUM MAPPING

We now wish to construct the conserved quantities associated with the obvious symmetry group. The technical tool by which this is done is called the momentum mapping. The reader not acquainted with this apparatus should consult the appendix.

First, let $\xi \in \mathfrak{g} = \mathfrak{so}(3)$. Next, solve the equation $\hat{dJ}(\xi) = \iota_{\xi_P} \omega$, where ξ_P is the infinitesimal generator of action corresponding to ξ .

To do this, we let ξ_1, ξ_2, ξ_3 be the usual basis for \mathfrak{g} . Then

$$(\xi_k)_P = -\varepsilon_k^{\ell} q^m \partial_{\ell} - \varepsilon_{k\ell}^m p_m \dot{\ell},$$

so that

$$\iota_{(\xi_k)_P} \omega = -\varepsilon_k^{\ell} q^m \omega_{\ell s} dq^s + d[\varepsilon_{km}^{\ell} q^m p_{\ell}].$$

We wish to express the first term as an exact differential. Some fiddling with indices yields

$$-\varepsilon_k^{\ell} q^m \omega_{\ell s} dq^s = \lambda d[q^k \|q\|^{-1}],$$

since $\omega_{\ell s} = \frac{1}{2} \frac{\lambda}{3} \varepsilon_{\ell st} q^t$, we may write.

$$\hat{J}(\xi_k) = \lambda q^k \|q\|^{-1} + \varepsilon_{km}^{\ell} q^m p_{\ell}, \quad (2)$$

and

$$\hat{J}(\xi) \cdot (q, p) = \langle \lambda \|q\|^{-1} q + q \times p, \xi_p \rangle.$$

Using the identification of $\mathfrak{so}(3)$ with (\mathbb{R}^3, \times) and \mathbb{R}^{3*} with \mathbb{R}^3 by $\langle \cdot, \cdot \rangle$, we get

$$J(q, p) \cdot \xi = \langle q \times p + \lambda \|q\|^{-1} q, \hat{\xi} \rangle, \quad (3)$$

which is the desired form of the momentum map.

Since $SO(3)$ is semi-simple, the co-adjoint cocycle associated to J vanishes and so J is Ad^* -equivariant. Thus \hat{J} is a homomorphism from \mathfrak{g} to the Lie algebra of functions under the Poisson bracket.

The derivative of J is

$$DJ_{(q,p)} \cdot (v, w) = \lambda \|q\|^{-1} [v - \|q\|^2 \langle q, v \rangle \cdot q] + q \times w + v \times p,$$

which tells us that all values of J are regular.

Furthermore, if $q \times p \neq 0$, then the isotropy group $G_\mu = SO(2)$ acts freely on $J^{-1}(\mu)$.

In this case, the conditions for reduction to work are satisfied and we may conclude that there is a two dimensional phase space P_μ with a unique symplectic form ω_μ . In other words,

$p_\mu = J^{-1}(\mu)/G_\mu$ has a unique ω_μ .

To examine the reduced system we work in polar coordinates, where the symplectic form is

$$\omega = \lambda \sin \phi \, d\phi \wedge d\theta + dr \wedge dp_r + d\theta \wedge dp_\theta + d\phi \wedge dp_\phi, \quad (4)$$

and the momentum integrals are

$$\begin{aligned} J^1 &= \lambda \cos \theta \sin \phi - \cos \theta \cot \phi \, p_\theta - \sin \theta \, p_\phi, \\ J^2 &= \lambda \sin \theta \sin \phi - \sin \theta \cot \phi \, p_\theta + \cos \theta \, p_\phi, \\ J^3 &= \lambda \cos \phi + p_\theta \end{aligned} \quad (5)$$

We reduce at $\mu = (0,0,L)$ and note that the following relations hold:

$$p_\phi = \cos \theta \, J^2 - \sin \theta \, J^1 = 0$$

$$p_\theta = \lambda \sin \phi \tan \phi.$$

These imply that

$$p_{\theta} = L - \lambda \cos \phi$$

$$= L(1 - \lambda^2 L^{-2}).$$

After a little algebra we get

$$h_{\mu} = \frac{1}{2m} \left((p_r)^2 + \frac{L^2}{r^2} \right) - \frac{k}{r}, \quad (6)$$

$$\omega_{\mu} = dr \wedge dp_r,$$

as the Hamiltonian structure on the reduced phase space

$$P_{\mu} = T^*R.$$

We note that the reduced system is identical to the reduced Kepler system.

3. THE HIDDEN SYMMETRY

A hidden symmetry is an infinitesimal symplectomorphism on P that

(HS1) is not an obvious symmetry,

(HS2) is tangent to an energy surface.

Since the set of all such vector fields forms an infinite dimensional Lie algebra, we will restrict ourselves to a geometrically meaningful finite dimensional subset which, in keeping with the approach for obvious symmetries, is again generated by complete lifts.

To construct such vector fields take the complete lift to P of a type $(1,1)$ tensor on Q and contract it with the second order equation. Add this vector field to the complete lift of a type $(1,0)$ tensor (a vector field) on Q to get a candidate for a generator of a hidden symmetry.

That is, let $f \in \mathcal{J}_1^1(Q)$ so that f has the local form $f = f_{\ell}^k \partial_k \otimes dq^{\ell}$. Then the complete lift of f , denoted f^C , has the property that $f^C \in \mathcal{J}_1^1(TQ)$, and has the local form

$$f^C = f_{\ell}^k \partial_k \otimes dq^{\ell} + f_{\ell}^k \partial_{\dot{k}} \otimes dv^{\ell} + v^k \partial_k f_{\ell}^m \partial_m \otimes dq^{\ell}. \quad (7)$$

The contraction then has the local form

$$\iota_{\chi_L} f^C = f_{\ell}^k v^{\ell} \partial_k + (v^j v^k \partial_j f_k^{\ell} + \xi^k f_k^{\ell}) \partial_{\ell} \quad , \quad (8)$$

where

$$\chi_L = v^k \partial_k + \xi^k \partial_{\dot{k}}$$

is the second order equation. We let T be a candidate to generate a hidden symmetry. That is,

$$T = \iota_{\chi_L} f^C + \tilde{\chi} \quad , \quad (9)$$

where $\tilde{\chi}$ is the complete lift of a vector field χ on Q . We now wish to solve the equations

$$L_T \omega = 0, \quad (10)$$

$$L_T E = 0,$$

where E is the energy.

With the usual methods of index gymnastics a straightforward, but long, calculation shows that the first equation implies that

$$f_{k\ell} = d_{k\ell m} q^m + e_{k\ell} \quad ,$$

where $f_{k\ell} = \delta_{km} f_{\ell}^m$,

$$d_{k\ell m} = d_{\ell km}, \quad (11)$$

$$e_{k\ell} = e_{\ell k},$$

and $d_{k\ell m} + d_{\ell mk} + d_{mk\ell} = 0.$

Letting $\chi = g^k \partial_k$ we find also that

$$\partial_{(k} g_{j)} + \partial_{(k} \xi^{\ell}_{\cdot} f_{j)\ell} = 0, \quad (12)$$

$$\partial_k [v^m \partial_m g_{\ell} + \partial_{\dot{\ell}} ((f_{mn} v^n + g_m) \xi^m)] = 0,$$

where again we have lowered indices with δ .

For the Kepler problem we know that we can solve these equations (11) and (12) with 3 different solutions

$$d_{jklm} = (\delta_{jm} \delta_{k\ell} - \delta_{j(k} \delta_{\ell)m}),$$

$$e_{jkl} = 0,$$

$$g_{jk} = 0,$$

where j acts like a tensor index. Using these as our starting point we find (after much manipulation) that we can again solve these for the monopole problem with the values

$$d_{jklm} = (\delta_{jm}\delta_{kl} - \delta_{j(k}\delta_{l)m}),$$

$$e_{jkl} = 0, \quad (13)$$

$$g_{jk} = \frac{\lambda}{r} \epsilon_{jklq} \ell.$$

If we let χ_{R^j} be the vector field formed with these values of d_{jklm} , e_{jkl} and g_{jk} , and find R^j which is defined by the relation

$$dR^j = i_{\chi_{R^j}} \omega,$$

we calculate that

$$R^j = m \epsilon_{km}^j v^k J^m - \frac{mk}{r} q^k, \quad (14)$$

and on the cotangent bundle that

$$R^j = \epsilon_{mp_k}^{jk} J^m - \frac{mk}{r} q^k. \quad (15)$$

From these it is straightforward to show that the Poisson brackets

satisfy

$$\begin{aligned}\{J^k, R^\ell\} &= \epsilon^{k\ell} R^m, \\ \{R^k, R^\ell\} &= (-2m\hbar) \epsilon^{k\ell} J^m.\end{aligned}\tag{16}$$

Before discussing these equations we first introduce some notation for constant energy surfaces:

$$\Sigma_e := \{x \in T^*Q \mid H(x) = e\}.$$

We also define the open submanifolds of P :

$$\Sigma_{\pm} := \bigcup_{\substack{e > 0 \\ e < 0}} \Sigma_e.$$

On Σ_{ϵ} we define

$$J_{\epsilon}^k := J^k|_{\Sigma_{\epsilon}},$$

$$K_{\epsilon}^k := (\epsilon 2m\hbar)^{-1/2} R^k|_{\Sigma_{\epsilon}},$$

$$\epsilon = \pm.$$

Now on Σ_{ϵ} we have the Poisson brackets:

$$\{J_{\varepsilon}^k J_{\varepsilon}^{\ell}\} = \varepsilon^{k\ell} J_{\varepsilon}^m ,$$

$$\{J_{\varepsilon}^k, K_{\varepsilon}^{\ell}\} = \varepsilon^{k\ell} K_{\varepsilon}^m , \quad (17)$$

$$\{K_{\varepsilon}^k, K_{\varepsilon}^{\ell}\} = -\varepsilon \varepsilon^{k\ell} J_{\varepsilon}^m .$$

So we see that under the Poisson bracket the functions L_{ε}^k and K_{ε}^{ℓ} generate the Lie algebras $so(3,1)$ and $so(4)$ on \sum_e for $\varepsilon = +$ and $\varepsilon = -$ respectively.

For the sequel we are only interested in the case \sum_- , so we abuse notation and remember that from now on J^k means J_{ε}^k and K^j means K_{ε}^j .

It is not difficult to show that the symplectic gradients of the J^k s and K^j s span an energy surface. This means that for any $e < 0$, $SO(4)$ has a symplectic action on P , which is transitive on \sum_e . In the next chapter we will try to realize this action.

4. Notes:

- (A) The obvious symmetry of the monopole was first discovered by Poincaré (1896) while explaining some seemingly paradoxical observations relating to experiments done on a Crooke's tube. In this paper Poincaré proved that, in the absence of a potential, an electrically charged particle has its motion confined to the surface of a cone, where the monopole is situated at the vertex. Furthermore, he showed that the particle motion is a geodesic, where the metric is the Euclidean metric induced by the imbedding of the cone in \mathbb{R}^3 .
- (B) This sort of result was investigated by Boulware et al (1976), who opened these cones out into a plane to discover that the motion on the cone corresponded to motion in an inverse square law potential in the corresponding plane. We call such a map an umbrella map, since the opening of the cone is similar to the way an umbrella is opened (we shall have more to say on this map later).
- (C) The hidden symmetry was first discovered by Zwanziger (1968) who introduced the extra term in the potential so that the Hamiltonian would look simpler in polar coordinates.

- (D) Schonfeld (1980) combined the extra potential term of Zwanziger and the umbrella map of Boulware to display the hidden symmetry. This works because the monopole problem is transformed into the Kepler problem.
- (E) The preceeding results, while very pretty, must be viewed as incomplete since none of the preceeding authors discuss the symplectic structure. Further, the umbrella map must be viewed as suspect from the viewpoint of symplectic geometry, since the umbrella map is not symplectic. In other words, when confronted with a non-canonical transformation we say our Hamiltonian creed: "What is canonical is important and what is important is canonical".
- (F) The cohomology class of ω shows us that there can be no symplectic map between the Kepler problem and the monopole problem. However, the reduction process shows us that it is possible for two inequivalent Hamiltonian systems to be acted on by the same group, with the same action, reduced at regular values and have the resulting reduced systems equivalent. Abraham and Marsden (1978) says that it is also possible to reduce exact symplectic manifolds and get non-exact symplectic forms on the reduced spaces. Thus it would seem that there are no cohomology theorems for momentum mappings.

- (G) Considering both (E) and (F), we would conclude that the geometric understanding of the umbrella map is on the level of the reduced systems, and not on the phase space level.
- (H) The construction of the hidden symmetry generators can be extended to $f \in \mathcal{F}_n^1(Q)$ for arbitrary positive n as follows:

Let $T = \iota_{\chi_L} \iota_{\chi_L} \dots \iota_{\chi_L} f^C$ where f^C is again the complete lift of f . That is, $f^C \in \mathcal{F}_n^1(TQ)$. Now T has the local coordinate form for f symmetric in the covariant part:

$$T = v^{k_1} \dots v^{k_n} f_{k_1 \dots k_n}^m \partial_m + (v^{k_1} \dots v^{k_n} v^\ell \partial_\ell f_{k_1 \dots k_n}^m + v^{k_1} \dots v^{k_{n-1}} \xi^{k_n} f_{k_1 \dots k_n}^m) \partial_m.$$

To see that such constructions are needed in mechanics, consider the n -dimensional Toda lattice, where a hidden symmetry exists as a polynomial of degree n for every n .

- (I) The outstanding problem remains, however. That is, to give an algorithmic understanding of when one can extend such procedures to generate finite dimensional Lie groups of

symmetries that include the obvious symmetry group. The situation is not all black however, for we see that if such a group is generated by vector fields that are the symplectic gradients of functions polynomial in the v^k 's, then such a construction is possible. Such an approach is also promising if one wishes to construct dynamical groups (see Marle (1976)).

(J) The reason for working on the tangent bundle instead of the cotangent bundle is essentially simplicity. As an example, compare the discussions of polynomial observables in Woodhouse (1980) and Kostant (1974). The essentials are not the complete story, for one has the freedom to construct lifts using either the canonical or the charged symplectic structure, and it is not a priori clear how this can be sorted out.

(K) The approach I have taken towards symmetries is due to the general philosophy that when one is working with geometric structures, one should only use constructions that depend on the given geometry. Concretely, the geometry given in the system has two facets: the symplectic structure of phase space and the metric structure of the individual cotangent spaces. I do not invoke the accidental Euclidean structure of phase space. I have endeavoured to provide a reasonable comprehensive account of the recent work in symmetries in

mechanics in general and applications to the Kepler problem in particular. The reader is invited to pursue the bibliography and note the different approaches taken. Lie transforms, Jacobi fields or Lagrangian dynamics (and others!) are represented. For instance, one can see Andrie and Simms (1972), Kunzle (1969), Levy-Leblond (1970), Mariwalla (1975), Rogers (1973) and Wolf (1977).

CHAPTER TWO

1. REALIZATION

This seems a suitable point to review our journey so far, and preview what is yet to come. In the προλεγόμενον we constructed the phase space, symplectic structure, and the Hamiltonian so that we knew precisely what mechanical system we were studying. Then, we set out to systematically find the symmetries of the system, both obvious and hidden. Now, we try to realize the symmetry. By this we mean that we try to find coordinates so that the symmetry now appears as more or less self-evident. Along the way, we also see how the symmetry gives a very elegant way to integrate the equations of motion.

Due to the algebraic and computational methods used in this section, we first transform to dimensionless variables to avoid excessive bulk in calculation and formulae.

For $k, \lambda > 0$ put

$$\vec{r} = \frac{km}{\lambda^2} \vec{q} \qquad r = \|\vec{r}\|$$

$$\vec{v} = \frac{\lambda}{mk} \vec{p} \qquad v = \|\vec{v}\|$$

$$\tau = \frac{mk^2}{\lambda^3} t \qquad \cdot = \frac{d}{dt} \quad .$$

We then have the relations

$$\vec{v} = \frac{d\vec{r}}{d\tau}$$

$$h = \frac{mk^2}{\lambda^2} H = \frac{1}{2} v^2 - \frac{1}{r} + \frac{1}{2r^2}$$

$$\vec{j} = \lambda \vec{J} = \vec{r} \times \vec{v} + \frac{1}{r} \vec{r}$$

$$\vec{k} = |k| \vec{R} = \vec{v} \times \vec{j} - \frac{1}{r} \vec{r}$$

$$\omega = \frac{1}{j} \omega_\lambda = \frac{1}{2} r^{-3} \varepsilon_{k\ell m} r^k dr^\ell dr^m + dr_k dv^k.$$

A short calculation gives

$$\vec{j} \cdot \vec{j} = r^2 v^2 - (\vec{r} \cdot \vec{v})^2 + 1$$

$$\vec{r} \cdot \vec{j} = r$$

$$\vec{r} \cdot \vec{k} = j^2 - r - 1$$

(A)

$$\vec{v} \cdot \vec{j} = \frac{1}{r} (\vec{r} \cdot \vec{v})$$

$$\vec{v} \cdot \vec{k} = -\frac{1}{r} (\vec{r} \cdot \vec{v})$$

$$\vec{j} \cdot \vec{k} = -1$$

$$\vec{k} \cdot \vec{k} = 2h(j^2 - 1) + 1.$$

The dimensionless form of the equations of motion is

$$\dot{\vec{r}} = \vec{v}$$

$$\dot{\vec{v}} = -\frac{1}{3} \frac{(\vec{r} \cdot \vec{j})}{r}.$$

Imitating Souriau (1974), we now define a new parameter s (the Levi-Civita time) by

$$s = \vec{r} \cdot \vec{v} - 2h\tau$$

so that

$$\frac{ds}{dt} = \frac{1}{r}; \text{ in other words } \frac{d}{ds} = r \frac{d}{dt}.$$

With the use of the previous formulae, we derive

$$\tau' = r \tag{1}$$

$$\tau'' = \vec{r} \cdot \vec{v} \tag{2}$$

$$\vec{r}' = r\vec{v} \tag{3}$$

$$\vec{r}'' = (\vec{r} \cdot \vec{v})\vec{v} - \frac{1}{r} \vec{r} + \frac{1}{r} \vec{j} \tag{4}$$

$$= (\vec{r} \cdot \vec{v})\vec{v} + \left(\frac{1}{2} - \frac{1}{r}\right)\vec{r} + \frac{1}{r} \vec{r} \times \vec{v} \tag{5}$$

where $'$ denotes d/ds .

Using (1), (2), and (4) we see

$$\vec{r}'' = \frac{1}{\tau'} [\tau'' \vec{r}' - \vec{r} + \vec{j}], \tag{6}$$

while differentiating (2) gives

$$\tau'' = 1 + 2h\tau' \tag{7}$$

Differentiating (6) and using (7) yields

$$\vec{r}'''' = 2h\vec{r}' \quad . \quad (8)$$

Now we call $\xi = \begin{pmatrix} \tau \\ \vec{r} \end{pmatrix}$, $\Xi^T = (\xi, \xi', \xi'', \xi''')$. The motivation is because now

$$\tau'''' = 2h\tau'' \quad (9)$$

$$\vec{r}'''' = 2h\vec{r}''$$

which, when translated into (ξ, Ξ) form, becomes the linear matrix differential system

$$\Xi' = A\Xi, \quad (10)$$

where

$$A = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 2h & 0 \end{pmatrix}.$$

The fundamental solution to the initial value problem (10) with

$$\Xi(s_0) = I_4 \quad \text{is}$$

$$\Xi(s) = e^{(s-s_0)A},$$

so we have reduced the integration of the equations of motion to exponentiating the matrix A .

Using the formulas (1) through (10) we can easily calculate the relations

$$\vec{r}''^2 - 2h\tau''^2 = 1 + 2h$$

$$\vec{r}''''^2 - 2h\tau''''^2 = -2h(1+2h)$$

$$\vec{r}'' \cdot \vec{r}'''' - 2h\tau''\tau'''' = 0$$

For $-\frac{1}{2} \leq h < 0$ we define ψ by $\psi^2 = -2h$, where ψ is positive.

Now define X and Y by

$$X = \begin{pmatrix} \psi \tau'' \\ \vec{r}'' \end{pmatrix} \quad Y = \begin{pmatrix} \tau'' \\ \frac{1}{\psi} \vec{r}'' \end{pmatrix} .$$

We see that X and Y satisfy

$$\|X\| = 1 - \psi^2$$

$$\|Y\| = 1 - \psi^2$$

$$X \cdot Y = 0 ,$$

so for $0 < \psi < 1$ an energy surface Σ_ψ has the topology of $S^3 \times S^2$. For $\psi = 1$ it is easy to see $\Sigma_{\psi=1} = S^2$. Summing this discussion up formally, we get the

THEOREM: (TOPOLOGY OF AN ENERGY SURFACE)

$$\Sigma_\psi = \begin{cases} S^3 \times S^2 & 0 < \psi < 1 \\ S^2 & \psi = 1 \end{cases} .$$

If we write

$$X = \begin{pmatrix} X^0 \\ \vec{X} \end{pmatrix} \quad Y = \begin{pmatrix} Y^0 \\ \vec{Y} \end{pmatrix}$$

we can invert the mapping

$$(\vec{r}, \vec{v}) \rightarrow ((\vec{X}, \vec{Y}), (X^0, Y^0, \psi))$$

to get

$$\vec{r} = \frac{1}{\psi^2 \mu^2} [(\mu^2 - Y^0) \vec{X} + X^0 \vec{Y} - \psi \vec{X} \times \vec{Y}]$$

$$\vec{v} = \frac{-\psi}{1 - Y^0} \vec{Y} .$$

We also calculate

$$\vec{j} = \frac{1}{\mu^2} [Y^0 \vec{X} - X^0 \vec{Y} + \frac{1}{\psi} \vec{X} \times \vec{Y}]$$

$$\vec{k} = \frac{-1}{\mu^2} [Y^0 \vec{X} - X^0 \vec{Y} + \psi \vec{X} \times \vec{Y}] ,$$

where $\mu^2 = 1 - \psi^2$.

With these variables we can easily describe $\sum_- = \bigcup_{0 < \psi \leq 1} \sum_\psi$. If we put

$$A = \begin{pmatrix} X \\ \psi \\ 0 \end{pmatrix} \quad B = \begin{pmatrix} Y \\ 0 \\ \frac{\psi}{j} \vec{j} \end{pmatrix}$$

then we have the relations

$$\|A\| = 1$$

$$\|B\| = 1$$

$$A \cdot B = 0.$$

Since \vec{j} can be described in terms of X and Y , we then may conclude the

THEOREM: (TOPOLOGY OF \sum_-)

$$\sum_- = S_+^4 \times S^2$$

where S_+^n is the open upper hemisphere of S^n .

These theorems propose the two natural questions:

- a) What is the symplectic structure on \sum_- ?
- b) How does $SO(4)$ act on \sum_- ?

to which we now turn.

When examining the symplectic form, one rapidly discovers that the variables X and Y are quite simply the wrong coordinates to use. An alternative is to examine the Poisson structure on some related manifold.

We define \vec{E} and \vec{F} by

$$\begin{aligned}\vec{E} &= \frac{1}{2}(\vec{j} + \frac{1}{\psi} \vec{k}) \\ \vec{F} &= \frac{1}{2}(\vec{j} - \frac{1}{\psi} \vec{k}).\end{aligned}$$

Then these satisfy the relations

$$\begin{aligned}\{E^k, E^\ell\} &= \epsilon^{k\ell}_m E^m \\ \{E^k, F^\ell\} &= 0 \\ \{F^k, F^\ell\} &= \epsilon^{k\ell}_m F^m.\end{aligned}\tag{11}$$

For notational convenience we also set

$$E = \|\vec{E}\| \quad \text{and} \quad F = \|\vec{F}\|.$$

Some calculation then shows

$$\begin{aligned}\vec{r} &= \frac{1}{A} [a\vec{E} + b\vec{F} + c\vec{E} \times \vec{F}] \\ \vec{v} &= \frac{1}{B} [-\chi F \vec{E} + E \chi \vec{F} - \xi \vec{E} \times \vec{F}]\end{aligned}\tag{12}$$

where

$$\begin{aligned}A &= (\psi^2(1+\psi)F(EF-G))^{-1} \\ a &= (F^2(\xi(1+\psi-\chi)+\chi^2)) \\ b &= [(EF-G)(\chi^2+\psi^2+\psi) - \chi^2G - EF\xi(1+\psi-\chi)] \\ c &= \chi F(\xi-(1+\psi-\chi)) \\ B &= (1-\xi)(EF-G)/\psi \\ G &= \vec{E} \cdot \vec{F} \\ \chi &= X^0 \\ \xi &= Y^0 \\ \psi &= 1/(E+F).\end{aligned}$$

We also have the constraints

$$F = 1 + E$$

$$\chi^2 + \xi^2 = 2(EF-G)/(E+F)^2.$$

We introduce a new variable σ by

$$\chi = \psi\sqrt{2(EF-G)} \cos \sigma$$

$$\xi = -\psi\sqrt{2(EF-G)} \sin \sigma$$

so we work in a seven dimensional space and derive that

$$\begin{aligned} \{E^k, \sigma\} = \cos^2 \sigma [& (\frac{a}{A} d - \frac{\chi F}{B} e + \psi f) E^k \\ & + (\frac{b}{A} d + \frac{\chi E}{B} e - \psi f) F^k \\ & + (\frac{c}{A} d - \frac{\xi}{B} e) (\vec{E} \times \vec{F})^k], \end{aligned}$$

$$\{F^k, \sigma\} = -\{E^k, \sigma\},$$

where

$$d = (\frac{1}{2y} + \frac{\xi}{2\chi^2} ((\frac{1-\xi}{y}) - (\frac{2y-1}{y^2})))$$

$$e = (\frac{\xi}{2\chi\psi} - \frac{y\psi}{2\chi})$$

$$f = (\frac{1}{2(1-\xi)} - \frac{\xi}{2\chi^2} (1 + \frac{1}{1-\xi}))$$

$$y = (E+F)(E+F - \sqrt{2(EF-G)} \sin \sigma).$$

From all this we see that $SO(4)$ acts as two independent copies of $SO(3)$ on \vec{E} and \vec{F} in the standard way, but in a highly convoluted way in σ . However, at last it is possible to 'see why' $SO(4)$ is in the magnetic problem, although the global topology can no longer be seen clearly.

Some concluding remarks are now in order. The presence of the magnetic field makes all calculations involving the symplectic form (or

dually, the Poisson structure) absolutely non-trivial. This is because the Poisson structure is

$$\{r^k, r^\ell\} = 0$$

$$\{r^k, v^\ell\} = \delta^{k\ell}$$

$$\{v^k, v^\ell\} = -r^{-3} \varepsilon_s^{k\ell} r^s.$$

One would hope that it would be possible to find a variable v such that the Poisson brackets $\{E^k, v\}$, $\{F^k, v\}$ had a very pleasing appearance. However, there is no a priori reason why such a variable should exist.

2. Notes:

- (A) The parameter s was first introduced by Levi-Civita (1906) to compactify the two-dimensional Kepler problem. The most systematic exploitation of s occurs in the work of Souriau. It is still completely amazing to me that it works so well in the monopole problem.
- (B) In the regularized Kepler problem all negative energy manifolds have the topology of $S^3 \times S^2$. The appearance of rest points is completely unique to the monopole.
- (C) For the variables (\vec{E}, \vec{F}, v) one can find 'nice' Poisson structures. However, the presence of $\vec{E} \cdot \vec{F}$ terms prevents us from actually realizing any of these structures concretely as the monopole problem.
- (D) One of the disappointing features of the monopole is the lack of a global symplectic potential. The 'disappointing' feature of this is that it prevents us from defining action-angle variables in any open set of configuration space. Weinstein has shown how to construct such variables for non-abelian groups (e.g. $SO(3)$), but it seems an open problem of how to generalize the construction to non-exact forms. Two resolutions seem possible. The first is to work on a $U(1)$ bundle over phase space where a global action exists. This seems somewhat unappetizing. The second is to

somehow understand the momentum map more fully. For example, if one pulls back the Kirillov-Kostant-Souriau symplectic form on the co-adjoint orbit of $SO(3)$ via the momentum map for the Kepler problem and subtracts it from the canonical symplectic form, one finds that the resulting presymplectic form is everywhere orthogonal to the angular momentum.

A.1 Notation and Defintions.

In this appendix we fix the notational conventions used in this thesis.

1. Polar Coordinates: The transformation from rectangular to polar coordinates is defined on an open subset of \mathbf{R}^3 by the map $T : (r, \theta, \phi) \rightarrow (q^1, q^2, q^3)$ given by

$$\begin{aligned} q^1 &= r \cos \theta \sin \phi \\ q^2 &= r \sin \theta \sin \phi \\ q^3 &= r \cos \phi \end{aligned}$$

Orthonormal (with respect to the standard inner product) bases for the tangent and cotangent spaces are given by

$$\begin{aligned} & \left(\frac{\partial}{\partial q^1}, \frac{\partial}{\partial q^2}, \frac{\partial}{\partial q^3} \right) \\ & (dq^1, dq^2, dq^3) . \end{aligned}$$

Coordinate bases in the polar system are given by

$$\begin{aligned} & \left(\frac{\partial}{\partial r}, \frac{\partial}{\partial \theta}, \frac{\partial}{\partial \phi} \right) \\ & (dr, d\theta, d\phi) . \end{aligned}$$

However, these are not orthonormal. Orthonormal, non-coordinate frames are given by

$$(\hat{\partial}_r, \hat{\partial}_\theta, \hat{\partial}_\phi)$$

$$(\hat{\theta}^r, \hat{\theta}^\theta, \hat{\theta}^\phi) .$$

These are related to the coordinate frame by the relations

$$\hat{\partial}_r = \frac{\partial}{\partial r}$$

$$\hat{\partial}_\theta = \frac{1}{r \sin \phi} \frac{\partial}{\partial \theta}$$

$$\hat{\partial}_\phi = \frac{1}{r} \frac{\partial}{\partial \phi}$$

$$\hat{\theta}^r = dr$$

$$\hat{\theta}^\theta = r \sin \phi d\theta$$

$$\hat{\theta}^\phi = r d\phi .$$

In the body of the thesis we make use of the relations

$$dq^1 \wedge dq^2 \wedge dq^3 = r^2 \sin \phi dr \wedge d\phi \wedge d\theta$$

$$\frac{1}{2} \varepsilon_{k\ell m} q^m dq^k \wedge dq^\ell = r^3 \sin \phi d\phi \wedge d\theta .$$

2. Invariant Scalars: The usual action of $SO(3)$ on R^3 induces an action on R^3 which leaves the functions

$$\begin{aligned} r &= (q^k q_k)^{1/2} \\ v &= (v^k v_k)^{1/2} \\ \alpha &= q^k v_k \end{aligned}$$

invariant. This notation is needed in the calculations on local equivalence.

3. Summation Convention: The summation convention of Einstein is used consistently unless followed by the sign for no sum: \nolimits .

4. Differentiation Conventions: Our notation for differentiation is defined by the relations

$$\partial_k f = \frac{\partial f}{\partial q^k}$$

$$\partial_{\dot{k}} f = \frac{\partial f}{\partial v^k}$$

$$\partial_{\dot{k}} f = \frac{\partial f}{\partial p_k} \quad .$$

Similar notation is used for vector fields:

$$X^k \partial_k + Y^\ell \partial_{\dot{\ell}} = X^k \frac{\partial}{\partial q^k} + Y^\ell \frac{\partial}{\partial v^\ell} \quad .$$

5. Background References: For background material on differential geometry see the books by Abraham-Marsden-Ratiu, Kobayashi-Nomizu, Poor, and Spivak. Material on characteristic classes can be found in Bott-Tu, Milnor-Stasheff, and Vaisman. For mechanics one should consult Abraham-Marsden, Arnold and Souriau. However, the best single reference in the spirit of this thesis is the book by Woodhouse on geometric quantization.

6. Notions from Differential Geometry

We consistently follow the notation of Abraham-Marsden (1978). For a manifold Q we let TQ denote its tangent bundle and let T^*Q denote its cotangent bundle. $\mathcal{T}_n^m(Q)$ denotes the tensors of type (m,n) on Q . For any tensor $T \in \mathcal{T}_n^m(Q)$ we may define a tensor $T^C \in \mathcal{T}_n^m(TQ)$ called the complete lift of T . We give local coordinate descriptions (which are easily checked to be invariant) for types $(1,0)$ and $(1,1)$ in the body of the thesis. For the definitive treatise on this branch of differential geometry see Yano and Ishihara (1973).

If G is a Lie group acting on a manifold Q and the Lie algebra of G is \mathfrak{g} , then the infinitesimal generator of action on Q (a vector field) corresponding to $\xi \in \mathfrak{g}$ is denoted by ξ_Q . Good references for Lie groups are Abraham-Marsden (1978), Cohn (1957), Chevalley (1946), and Helgason (1978).

7. The Momentum Mapping.

Let (P, ω) be a connected symplectic manifold and $\Phi : G \times P \rightarrow P$ a symplectic action of the Lie group G on P . By this we mean that for each $a \in G$, $\Phi_a^* \omega = \omega$. Let G have Lie algebra \mathfrak{g} and dual \mathfrak{g}^* . We then call the map $J : P \rightarrow \mathfrak{g}^*$ a momentum mapping for the action Φ if for each $\xi \in \mathfrak{g}$

$$dJ(\xi) = i_{\xi_P} \omega$$

where $\hat{J}(\xi) : P \rightarrow \mathbb{R}$ is defined by $\hat{J}(\xi)(x) = J(x) \cdot \xi$. The important thing to remember is that this is simply the generalization of the geometric ideas behind translational and rotational invariance giving us linear and angular momentum. For more details see Abraham-Marsden (1978), Souriau (1970), Woodhouse (1980), Cushman (1974) and Atiyah-Bott (1984).

BIBLIOGRAPHY

- Abraham, R., and Marsden, J. 1978. Foundations of Mechanics. Benjamin-Cummings, Reading, Mass.
- Abraham, R., Marsden, J., and Ratiu, T. 1982. Manifolds, Tensor Analysis, and Applications. Addison-Wesley, Reading, Mass.
- Aldaya, A. and de Azcarraga, J. 1982. Quantization as a consequence of the symmetry group: an approach to geometric quantization. J. Math. Phys. 7:1297-1305.
- Anderson, R., and Davison, S. 1974. A generalization of Lie's counting theorem for second-order ordinary differential equations. J. Math. Anal. and Appl. 48:301-315.
- Andrie, M., and Simms, D. 1972. Constants of motion and Lie group actions. J. Math. Phys. 3:331-336.
- Arens, R. 1964. Differential-geometric elements of analytic dynamics. J. Math. Anal. Appl. 9:165-202.
- Arnold, V. I.
1967. Characteristic class entering in quantization conditions. Funct. Anal. Appl. 1(1):1-13.
1973. Ordinary differential equations. MIT, Cambridge, Mass.
1978. Mathematical methods of classical mechanics. MIR (Moscow, 1975), Springer Graduate Texts in Math. No.60 Springer-Verlag, New York.
1983. Geometrical methods in the theory of ordinary differential equations. Springer-Verlag, New York.
- Atiyah, M., and Bott, R. 1984. The moment map and equivariant cohomology. Top. 1:1-28.
- Avez, A., Lichnerowicz, A., and Diaz-Miranda, A., 1974. Sur l'algebre des automorphismes infinitesimaux d'une variete symplectique. J. Diff. Geom. 9:1-40.
- Bacry, H. 1966. The de Sitter Group $L_{4,1}$ and the bound states of the hydrogen atom. Nuovo Cimento 41A:222-234.
- Bacry, H., Ruegg, H., and Souriau, J. M. 1966. Dynamical groups and spherical potentials in classical mechanics. Commun. Math. Phys. 3:323-333.
- Balachandran, A., Marmo, G., Skagerstam, B., and Stern, A.
1980. Magnetic monopoles without strings. Nucl Phys. B162:385-396.
1980. Supersymmetric point particles and monopoles with no strings. Nucl. Phys. B164:427-444.

- Barut, A., and Bornzin, G. 1971. $SO(4,2)$ formulation of the symmetry breaking in relativistic Kepler problems with or without magnetic charges. *J. Math. Phys.* 5:841-846.
- Bander, M., and Itzykson, C. 1966. Group theory and the hydrogen atom (I). *Rev. Mod. Phys.* 2:330-345.
- Birkhoff, G. D.
 1922. Surface transformations and their dynamical applications. *Acta. Math.* 43:1-119.
 1966. *Dynamical Systems*. Colloq. Publ. IX, 2nd ed. Am. Math. Soc., Providence, R.I.
- Bott, R., and Tu, L. 1982. *Differential forms in algebraic topology*. Springer, New York.
- Boulware, D., Brown, L., Cahn, R., Ellis, S., and Lee, C. 1976. Scattering on magnetic charge. *Phys. Rev. D.* 10:2708-2727.
- Cant. A. 1981. Invariant connections and magnetic monopoles. *J. Math. Phys.* 22:2283-2288.
- Caratu, G., Marmo, G., Saletan, E. J., Simoni, A., and Vitali, B. 1974. Homogeneous canonical transformations in phase space. *Nouvo Cimento* 19B:228-238.
- Carrigan, R. 1965. Consequences of the existence of massive magnetic poles. *Nuovo Cimento.* 1:638-641.
- Caviglia, G. 1983. Dynamical symmetries: an approach to Jacobi fields and to constants of geodesic motion. *J. Math. Phys.* 8:2065-2069.
- Chevalley, C. 1946. *Theory of Lie groups*. Princeton Univ. Press, Princeton, N.J.
- Choquet-Bruhat, Y., De Witt-Morette, C., and Dillard-Bleick, M. 1977. *Analysis, manifolds and physics*. North-Holland, New York.
- Cohn, P. M. 1957. *Lie groups*. Cambridge Univ. Press. Cambridge.
- Crampin, M. 1981. On the differential geometry of the Euler-Lagrange equations, and the inverse problem of Lagrangian dynamics. *J. Phys. A.* 14:2567-2575.
- Cushman, R. 1974. The momentum mapping of the harmonic oscillator, in *Inst. Nat. di Alta Mate., Symp. Math.* XIV:323-342. Academic, London.
- Czyz, J. 1979. On geometric quantization and its connections with the Maslov theory. *Rep. Math. Phys.* 1:57-97.

- D'Hoker, E., and Vinet, L.
 1983. Supersymmetry of the Pauli equation in the presence of a magnetic monopole. Preprint.
 1984. Dynamical supersymmetry of the magnetic monopole and the $1/r^2$ -potential. Preprint.
 1984. Superspace formulation of the dynamical symmetries of the Dirac magnetic monopole. Preprint.
 1984. A system with dynamical supersymmetry: the electron in the field of the magnetic monopole. To appear.
- Dickson, L. 1924. Differential equations from the group standpoint. *Ann. of Math.* 25:287-378.
- Dirac, P. A. M.
 1931. Quantized singularities in the electromagnetic field. *Proc. R. Soc. London* A133:60-72.
 1950. Generalized Hamiltonian dynamics. *Can. J. Math.* 2:129-148.
- Djukic, D. D. 1974. Conservation laws in classical mechanics for quasi-coordinates. *Arch. Rat. Mech. Anal.* 56:79-98.
- Dombrovski, P. 1962. On the geometry of the tangent bundle. *J. Reine Angew. Math.* 210:73-88.
- Dongpei, Z. 1981. Noether symmetry of the single-particle system. *J. Phys. A* 14:2807-2816.
- Duval, C. 1982. Nouvelles symetries de la mecanique newtonienne. Thesis, Univ. of Aix-Marseille.
- Duval, C., and Horvathy, P. 1982. Particles with internal structure: the geometry of classical motions and conservation laws. *Ann. of Phys.* 1:10-33.
- Estabrook, F. B., and Wahlquist, H. D. 1975. The geometric approach to sets of ordinary differential equations and Hamiltonian dynamics. *SIAM Review* 17(2):201-220.
- Flanders, H. 1963. Differential forms. Academic, New York.
- Fleming, W. 1965. Functions of several variables. Addison-Wesley, Reading, Mass.
- Fradkin, D. 1967. Existence of the dynamic symmetries O_4 and SU_3 for all classical central potential problems. *Prog. Theor. Phys.* 5:798-812.
- Giachetti, R. 1981. Hamiltonian systems with symmetry: an introduction. *Rivista del nuovo cimento* 4:12.
- Goldstein, H. 1950. Classical mechanics. Addison-Wesley, Reading, Mass.

- Gonzalez-Gascon, F. 1977. Notes on the symmetries of systems of differential equations. *J. Math. Phys.* 9:1763-1767.
- Gotay, M. J., and Nester, J. E. 1980. Generalized constraint algorithm and special presymplectic manifolds, in *Geometric methods in mathematical physics*, Kaiser, G., and Marsden, J., (eds.). Springer lec. notes in math. no. 775. Springer, New York.
- Gotay, M. J., Nester, J. E., and Hinds, G. 1978. Presymplectic manifolds and the Dirac-Bergman theory of constraints. *J. Math. Phys.* 19:2388-2399.
- Guillemin, V., and Sternberg, S.
 1977. *Geometric asymptotics*. Am. Math. Soc. Survey, vol. 14.
 1978. On the equations of motion of a classical particle in a Yang-Mills field and the principle of general covariance. Hadronic, Nonantum, Mass.
- Gyorgyi, G.
 1968. Kepler's equation, Fock variables, Bacry's generators and Dirac brackets. *Nuovo Cimento* 53A:717-735.
 1969. Kepler's equation, Fock variables, Bacry's generators II. *Nuovo Cimento* 62A:449-473.
- Havas, P. 1957. The range of application of the Lagrange formalism. *Nuovo Cimento* 3:363-388.
- Helgason, S. 1978. *Differential geometry, lie groups, and symmetric spaces*. Academic, New York.
- Hermann, R.
 1966. *Lie groups for physicists*. Benjamin-Cummings, Reading, Mass.
 1972. Spectrum-generating algebras in classical mechanics. I, II. *J. Math. Phys.* 13:833, 878.
- Hill, E. 1951. Hamilton's Principle and the conservation theorems of mathematical physics. *Rev. Mod. Phys* 3:253-260.
- Hirsch, M. W., and Smale, S. 1974. *Differential equations, dynamical systems and linear algebra*. Academic, New York.
- Hitchin, N. 1982. *The geometry of monopoles*. Lecture notes.
- Hoffman, K., and Kunze, R. 1961. *Linear algebra*. Prentice-Hall, Englewood Cliffs, N. J.
- Ikeda, M., and Maekawa, T. 1970. On finite transformations generated by the Runge-Lenz vector in the n-dimensional Kepler problem.

- Iwai, T.
 1980. On reduction of the four-dimensional harmonic oscillator. *J. Math. Phys.* 8:1628-1632.
 1980. On a "conformal" Kepler problem and its reduction. *J. Math. Phys.* 8:1633-1639.
- Jaffe, A., and Taubes, C. 1980. *Vortices and monopoles.* Birkhauser, Boston.
- Jost, R. 1964. Poisson brackets (an unpedagogical lecture). *Rev. Mod. Phys.* 36:572-579.
- Kaplan, W. 1942. Topology of the two-body problem. *Am. Math. Monthly* 49:316-323.
- Kobayashi, S. 1973. *Transformation groups in Riemannian geometry.* Springer, New York.
- Kobayashi, S., and Nomizu, K. 1963. *Foundations of differential geometry.* Wiley, New York.
- Kostant, B. 1974. Symplectic spinors, in *Conv. di geom. simp. e fisica math. Symp. Math.* XIV:139-152.
- Kummer, M. 1982. On the regularization of the Kepler problem. *Comm. Math. Phys.* 84:133-152.
- Kunzle, H.
 1969. Degenerate Lagrangian systems. *Ann. Inst. H. Poincare (A)*. 11:393-414.
 1972. *Ann. Inst. H. Poincare* 27. 337-362.
- Lanczos, C. 1962. *The variational principles of mechanics*, 4th ed. Univ. of Toronto Press.
- Levi-Civita, T.
 1906. Sur la resolution qualitative du probleme restreint des trois corps. *Acta Math.* 30:305-327.
 1906. Sur la regularisation du probleme des trois corps. *Acta Math.* 30:49-92.
- Levy-Leblond, J. M.
 1969. Group theoretical foundations of classical mechanics; the Lagrangian gauge problem. *Comm. Math. Phys.* 12:64-79.
 1970. Conservation laws for gauge-variant Lagrangians in classical mechanics. *Am. J. Phys.* 39:502-506.
- Ligon, T., and Schaaf, M. 1976. On the global symmetry of the classical Kepler problem. *Rep. on Math. Phys.* 3:281-300.
- Loomis, L. H., and Sternberg, S. 1968. *Advanced calculus.* Addison-Wesley, Reading, Mass.

- Lovelock, D., and Rund, H. 1975. Tensors, differential forms, and variational principles. Wiley, New York.
- Lutzky, M. 1977. Symmetry groups and conserved quantities for the harmonic oscillator. *J. Phys. A* 2:249-258.
- MacLane, S. 1970. Hamiltonian mechanics and geometry. *Am. Math. Monthly* 77:570-586.
- Mariwalla, K. H.
 1975. Dynamical symmetries in mechanics. *Phys. Rep. C* 5:287-362.
 1982. Integrals and symmetries: the Bernoulli-Laplace-Lenz vector. *J. Phys. A* 9:L467-L471.
- Marle, G. M. 1976. Symplectic manifolds, dynamical groups and Hamiltonian mechanics, in *Differential geometry and relativity*. M. Cahen and M. Flato, (eds.). D. Reidel.
- Marmo, G., and Saletan, E. J. 1977. Ambiguities in Lagrange and Hamiltonian formalism: transformation properties. *Nuovo Cimento* 40:67-89.
- Marmo, G., Saletan, E. J., and Simon, A. 1979. Reduction of symplectic manifolds through constants of the motion. *Nuovo Cimento* 50B:21-36.
- Marsden, J. E.
 1968. Generalized Hamiltonian mechanics. *Arch. Rat. Mech. Anal.* 28:326-362.
 1974. *Applications of global analysis in mathematical physics*. Publish or Perish, Boston.
 1981. *Lectures on geometric methods in mathematical physics*. SIAM, Philadelphia.
- Marsden, J. and Weinstein, A. 1974. Reduction of symplectic manifolds with symmetry. *Rep. Math. Phys.* 5:121-130.
- Meinhardt, J. 1981. Symmetries and differential equations. *J. Phys. A* 14:1893-1914.
- Menzio, M. R., and Tulczyjew, W. M. 1978. Infinitesimal symplectic relations and generalized Hamiltonian dynamics. *Ann. Inst. H. Poincare* 28A 349-367.
- Meyer, K. R. 1973. Symmetries and integrals in mechanics, in Peixoto, ed., *Dynamical systems*. pp. 259-273. Academic, New York.
- Miller, J. G. 1976. Charge quantization and canonical quantization. *J. Math. Phys.* 17:643-646.
- Milnor, J. 1983. On the geometry of the Kepler problem. *Am. Math. Monthly* 90:353-365.

- Milnor, J., and Stasheff, J. 1974. Characteristic classes.
Ann. of math. no. 76. Princeton univ. press. Princeton, N.J.
- Montgomery, R. 1983. Canonical formulations of a classic particle in a Yang-Mills field and Wong's equations. Preprint.
- Moriyasu, K.
1978. Gauge invariance rediscovered. Am. J. Phys. 3:274-278.
1981. Gauge invariance and hidden symmetries. Am. J. Phys. 9:819-826.
- Moser, J.
1958. New aspects on the theory of Hamiltonian systems.
Comm. Pure Appl. Math. 9:81-114.
1968. Lectures on Hamiltonian systems, in Memoirs Am. Math. Soc., vol. 81. Am. Math. Soc., Providence, R.I.
1970. Regularization of Kepler's problem and the averaging method on a manifold. Comm. Pure Appl. Math. 23.
- Muncaster, R. 1982. An abstract approach to group invariance, with applications to differential equations. Arch. for Rat. Mech. and Anal. 78:1-28.
- Neboroshev, N. 1972. Action-angle coordinates and their generalizations. Trud. Mosk. Math. 26.
- Onofri, E. 1976. Dynamical quantization for the Kepler manifold. J. Math. Phys. 3:401-408.
- Onofri, E., and Pauri, M.
1972. Dynamical quantization. J. Math. Phys. 13(4):533-543.
1973. Constants of the motion and degeneration in Hamiltonian systems. J. Math. Phys. 14:1106-1115.
- Pars, L. 1965. A treatise on analytical dynamics. Wiley, New York.
- Pin, Ong Chong, 1975. Curvature and mechanics. Adv. Math 15(3):269-311.
- Poincare, H.
1892. Les methodes nouvelles de la mecanique celeste, 1, 2, 3. Gauthier-Villars, Paris (1892); Dover, New York, 1957.
1896. Remarques sur une experience de M. Birkeland. C. R. Acad. Sci. Paris B123:530-533.
- Poor, W. 1981. Differential geometric structures. McGraw-Hill, New York.
- Prince, G., and Eliezer, C.
1980. Symmetries of the time-dependent N-dimensional oscillator. J. Phys. A 13:815-823.
1981. On the Lie symmetries of the classical Kepler problem. J. Phys. A 14:587-596.

- Roels, J., and Weinstein, A. 1971. On functions whose Poisson brackets are constant. *J. Math. Phys.* 12:1482-1486.
- Rogers, H. 1973. Symmetry transformations of the classical Kepler problem. *J. Math. Phys.* 8:1125-1129.
- Sanders, P. 1966. Magnetic charge. *Cont. Phys.* 6:419-429.
- Schafir, R. 1981. On the Lenz vector. *J. Phys. A*:14:L269-L271.
- Schonfeld, J. 1980. Dynamical symmetry and magnetic charge. *J. Math. Phys.* 10:2528-2533.
- Simms, D. 1974. Geometric quantization of energy levels in the Kepler problem. *Symp. Math.* XIV:125-137.
- Simms, D. J., and Woodhouse, N. M. J. 1976. Lectures on geometric quantization, *Lec. Notes Phys.*, vol. 53. Springer, New York.
- Smale, S. 1970. Topology and mechanics (I,II). *Inv. Math.* 10:305-331; 11:45-64.
- Sniatycki, J.
 1973. Prequantization of charge. *J. Math. Phys.* 5:619-620.
 1978. Geometric quantization and quantum mechanics. Springer, New York.
- Sommers, P. 1973. On Killing tensors and constants of motion. *J. Math. Phys.* 6:787-790.
- Souriau, J. M.
 1970. Structure des systemes dynamiques. Dunod, Paris.
 1974. Sur la variete de Kepler. *Symp. Math.* XIV:343-360.
 1975. Geometrie symplectique et physique mathematique, in *Coll. Intern. du CNRS*, no. 237, Paris.
 1982. Geometrie globale du probleme a deux corps. preprint.
- Spivak, M.
 1965. Calculus on manifolds. Benjamin-Cummings, Reading, Mass.
 1974. A comprehensive introduction to differential geometry (five vols). Publish or Perish, Boston.
- Sternberg, S.
 1963. Lectures on differential geometry. Prentice-Hall. Englewood Cliffs, N. J.
 1977. On Minimal coupling and the symplectic mechanics of a classical particle in the presence of a Yang-Mills field. *Proc. Nat. Acad. Sci.* 74:5253-5254.
- Sternberg, S., and Ungar, T. 1978. Classical and prequantized mechanics without Lagrangians or Hamiltonians. *Hadronic J.* 33-76.

- Synge, J. L.
 1926. On the geometry of dynamics. *Phil Trans. A* 220:31-106.
 1960. Classical dynamics. *Encycl. Phys.* 3:1-25.
- Synge, J. L., and Griffith, B. A. 1959. *Principles of mechanics*. McGraw-Hill, New York.
- T'Hooft, G. 1974. Magnetic monopoles in unified gauge theories. *Nucl. Phys.* B79:276-284.
- Tripathy, K. C. 1976. Dynamical prequantization, spectrum generating algebras and the classical Kepler and harmonic oscillator problems, in 1975 Fourth Intern. Coll. on Group Theor. Methods in phys., Nijmegen. *Lec. Notes in Phys.* vol. 50, pp. 199-209. Springer, New York.
- Tripathy, K., Gupta, R., and Anand, J. 1975. $O(4,2)$ symmetry and the classical Kepler problem. *J. Math. Phys.* 5:1139-1144.
- Tulczjew, W. M.
 1974. Hamiltonian systems, Lagrangian systems and the Legendre transformation. *Symp. Math.* XIV:247-258.
 1977. The Legendre transformation. *Ann. Inst. H. Poincare* 27:101-114.
- Vaisman, I. 1973. *Cohomology and differential forms*. Marcel Decker. New York.
- Vinogradov, A., and Kupershmidt, B. 1977. The structures of Hamiltonian mechanics. *Russ. Math. Surveys* 32:177-243.
- Wallach, N. 1977. *Symplectic geometry and Fourier analysis*. Math Sci, Brookline, Mass.
- Weinstein, A.
 1977. Lectures on symplectic manifolds, in C. B. M. S. Conf. Series., Am. Math. Soc., no. 29. Am. Math. Soc., Providence, R.I.
 1978. A universal phase space for particles in Yang-Mills fields. *Lett. Math. Phys.* 2:417-420.
 1983. The local structure of Poisson manifolds. *J. Diff. Geom.* 18:523-557.
- Wolf, J. 1977. Representations associated to minimal co-adjoint orbits, in differential geometric methods in mathematical physics II (proc. Bonn 1977), Springer lec. notes in math. vol. 676, 1978. Springer, New York.
- Wu, T., and Yang, C.
 1975. Concept of non-integrable phase factors and global formulation of gauge fields. *Phys. Rev. D* 12:3845-3857.
 1976. Dirac monopole without strings: monopole harmonics. *Nucl. Phys.* B107:365-380.

- Westenholz, C. Von. 1981. Differential forms in mathematical physics. North-Holland, Amsterdam.
- Whittaker, E. T. 1959. A treatise on the analytical dynamics of particles and rigid bodies, 4th ed. Cambridge Univ. Press, Cambridge.
- Woodhouse, N. 1980. Geometric quantization. Clarendon, Oxford.
- Wulfman, C., and Wyburne, B. 1976. The Lie group of Newton's and Lagrange's equations for the harmonic oscillator. J. Phys. A:4:507-518.
- Yano, K., and Ishihara, S. 1973. Tangent and cotangent bundles. Marcel Dekker, New York.
- Zwanziger, D. 1968. Exactly soluble nonrelativistic model of particles with both electric and magnetic charges. Phys. Rev. 5:1480-1488.

B30411